



© 2017, Marley, C.J., Hodson, D., Brugniaux, J.V., Fall, L. and Bailey, D.M.
This is an author produced version of a paper published in publication in
CLINICAL SCIENCE. Uploaded in accordance with the publisher's self-
archiving policy.

Marley, C.J., Hodson, D., Brugniaux, J.V., Fall, L. and Bailey,
D.M., 2017. Post-prandial hyperlipidaemia results in systemic
nitrosative stress and impaired cerebrovascular function in the
aged. *Clinical Science*, p.CS20171406.

Post-prandial hyperlipidaemia results in systemic nitrosative stress and impaired cerebrovascular function in the aged

Christopher J. Marley, PhD¹; Danielle Hodson, PhD^{1,2}; Julien V. Brugniaux^{1,3}, PhD; Lewis Fall, PhD¹ and Damian M. Bailey, PhD, FPVRI, FRSC, FACSM, FPS¹

¹Neurovascular Research Laboratory, Faculty of Life Sciences and Education, University of South Wales, UK; ²School of Social and Health Sciences, Leeds Trinity University, UK and ³School of Science and Health, Western Sydney University, Australia

Correspondence:

Professor Damian M. Bailey, Neurovascular Research Laboratory, Faculty of Life Sciences and Education, University of South Wales, UK, CF37 4AT.

Tel: +44 (0)1443 652296

Fax: +44 (0)1443 652285

e-mail: damian.bailey@southwales.ac.uk

Running Title: Post-prandial oxidative-nitrosative stress and cerebrovascular dysfunction

Key words: Aging, Cerebrovascular Function, High-Fat Meal, Hyperlipidaemia, Oxidative-Nitrosative Stress.

1 Abstract

2 Post-prandial hyperlipidaemia (PPH) acutely impairs systemic vascular endothelial function,
3 potentially attributable to a free radical-mediated reduction in vascular nitric oxide (NO)
4 bioavailability (oxidative-nitrosative stress). However, it remains to be determined whether
5 this extends to the cerebrovasculature. To examine this, 38 (19 young (≤ 35 years) and 19 aged
6 (≥ 60 years)) healthy males were recruited. Cerebrovascular function (middle cerebral artery
7 velocity, MCAv) and cerebrovascular reactivity to hypercapnea ($\text{CVR}_{\text{CO}_2\text{Hyper}}$) and hypocapnea
8 ($\text{CVR}_{\text{CO}_2\text{Hypo}}$) were determined via trans-cranial Doppler ultrasound and capnography. Venous
9 blood samples were obtained for the assessment of triglycerides (photometry), glucose
10 (photometry), insulin (radio-immunoassay), ascorbate free radical ($\text{A}^{\cdot-}$, electron paramagnetic
11 resonance spectroscopy) and nitrite (NO_2^- , ozone-based chemiluminescence) in the fasted state
12 prior to and 4 hours following consumption of a standardised high-fat meal (1,362 kcal; 130g
13 of fat). Circulating triglycerides, glucose and insulin increased in both groups following the
14 high-fat meal ($P < 0.05$), with triglycerides increasing by 1.37 ± 1.09 mmol/L in the young and
15 1.54 ± 1.00 mmol/L in the aged ($P < 0.05$). This resulted in an increased systemic formation
16 of free radicals in the young ($P < 0.05$) but not the aged ($P > 0.05$) and corresponding reduction
17 in NO_2^- in both groups ($P < 0.05$). While the meal had no effect on MCAv in either age group,
18 $\text{CVR}_{\text{CO}_2\text{Hyper}}$ was selectively impaired in the aged ($P < 0.05$). These findings indicate that PPH
19 causes acute cerebrovascular dysfunction in the aged subsequent to systemic nitrosative stress.

20

21

22

23

24

25 **Abbreviations:** $A^{\bullet-}$, ascorbate free radical; BP, blood pressure; BMI, body mass index;
26 CVR_{CO_2} , cerebrovascular reactivity to changes in CO_2 ; CVR_{CO_2Hyper} , cerebrovascular reactivity
27 in response to hypercapnea; CVR_{CO_2Hypo} , cerebrovascular reactivity in response to hypocapnea;
28 CVs, coefficient of variations; EPR, electron paramagnetic resonance; MAP, mean arterial
29 pressure; MCAv, middle cerebral artery velocity; NO, nitric oxide; NO_2^- , nitrite; PET_{CO_2} ,
30 partial pressure of end-tidal CO_2 ; PPH, post-prandial hyperlipidaemia.

31 Introduction

32 The consumption of a high-fat meal is characterised by a state of post-prandial hyperlipidaemia
33 (PPH) that can last for as long as 8 hours (1). During this time course, PPH has been shown to
34 acutely impair systemic vascular endothelial function (2-5), that may be related to a free
35 radical-mediated reduction in the vascular bioavailability of nitric oxide (NO) referred to as
36 oxidative-nitrosative stress (6). However, the extent to which PPH impacts the cerebral vessels
37 and may contribute to stroke risk has yet to be investigated. The reactivity of the cerebral
38 vessels in response to changes in carbon dioxide (CVR_{CO_2}) may be particularly vulnerable to
39 PPH, since it is considered a reflection of cerebrovascular endothelial function (7). This would
40 be of concern since any impairment in CVR_{CO_2} translates into an increased risk of stroke (8).
41 To date, only one study has investigated the impact of PPH on the cerebrovasculature and
42 reported that it had no effect on young adults (9). Whether this is the same in the more
43 “vulnerable, redox-reactive” aged brain remains to be established. To examine this, we
44 designed a study with young and aged adults to determine if PPH in the latter would be
45 associated with more marked impairments in cerebrovascular function subsequent to an age-
46 related increase in systemic oxidative-nitrosative stress.

47

48 **Materials and Methods**

49 *Participants*

50 Following ethical approval (#12/11/2013) and written informed consent, 19 young males (aged
51 25 ± 6 years with a body mass index (BMI) of 23 ± 4 kg.m²) and 19 aged males (aged 67 ± 5
52 years, BMI of 27 ± 3 kg.m²) were recruited. All participants were non-smokers, without a
53 history of cardiovascular, cerebrovascular or respiratory disease. All participants were
54 encouraged to follow a low nitrate/nitrite diet and abstain from any antioxidant supplements
55 prior to and throughout the duration of the study with specific instructions to avoid fruits, salads
56 and cured meats.

57

58 *Experimental Design*

59 Data collection began at 08:00 hours following a 12-hour overnight fast. Measurements of
60 cerebrovascular function and blood samples were obtained from each participant prior to and
61 4 hours following consumption of a standardised high-fat meal. The 4-hour mark was chosen
62 as this is when the triglyceride response is considered to be at its peak (1). All procedures
63 followed were in accordance with institutional guidelines.

64

65 *High-Fat Meal*

66 The test meal comprised of 130g of fat, 48g of carbohydrate and 9.5g of protein, as previously
67 described by Patsch and colleagues (1). The total energy content of the meal was 1,362 calories.

68

69

70

71 *Experimental Procedures*

72 *Cerebrovascular Function*

73 Middle cerebral artery velocity (MCAv) was determined using a 2MHz pulsed Doppler
 74 ultrasound system (Multi-Dop X4, DWL Elektronische Systeme GmbH, Sipplingen, Germany).
 75 Systolic, diastolic and mean arterial blood pressure (MAP) were monitored using finger
 76 photoplethysmography (Finometer PRO, Finapres Medical Systems, Amsterdam, The
 77 Netherlands), with pulse pressure calculated as systolic minus diastolic blood pressure.
 78 Changes in the partial pressure of end-tidal CO₂ (PET_{CO2}) were determined via capnography
 79 (ML 206, ADInstruments Ltd, Oxford, UK). Data were sampled continuously at 1kHz and
 80 stored for off-line analysis. Following 10 minutes breathing room air, CVR_{CO2} was assessed in
 81 response to hypercapnea (CVR_{CO2Hyper}) by breathing an inspirate of 5% CO₂ with 21% O₂ and
 82 balanced nitrogen for 3 minutes. Following a 5-minute recovery breathing room air, CVR_{CO2}
 83 was also assessed in response to hypocapnea (CVR_{CO2Hypo}) via 3 minutes of controlled
 84 hyperventilation (15 breaths.min⁻¹). Both CVR_{CO2Hyper} and CVR_{CO2Hypo} were calculated as the
 85 percentage change in MCAv from baseline per 1 mmHg change in PET_{CO2} recorded during the
 86 final 30 seconds (average taken) of the respective challenge when steady-state had been
 87 achieved:

88

$$89 \text{ CVR}_{\text{CO}_2} (\% \cdot \text{mmHg}^{-1}) = 100 \times \frac{\Delta \text{MCAv} (\%)}{\Delta \text{PET}_{\text{CO}_2} (\text{mmHg})} / [\text{PET}_{\text{CO}_2} (\text{final}) - \text{PET}_{\text{CO}_2} (\text{baseline})]$$

90

91 From these data, we derived the CVR_{CO2} range as a useful indication of the cerebral
 92 circulation's combined ability to respond to differential changes in CO₂ (10). This was
 93 calculated as the sum of the fractional vasodilation and vasoconstriction incurred during the
 94 respective hypercapnea and hypocapnea challenges as described:

95

$$96 \text{ CVR}_{\text{CO}_2} \text{ range } (\% \cdot \text{mmHg}^{-1}) = \text{CVR}_{\text{CO}_2\text{Hyper}} (\% \cdot \text{mmHg}^{-1}) + \text{CVR}_{\text{CO}_2\text{Hypo}} (\% \cdot \text{mmHg}^{-1})$$

97 ***Blood Samples***

98 Venous blood samples were drawn without stasis from a forearm antecubital vein via an 18-
99 gauge cannula into vacutainers. Once obtained, samples were centrifuged at 600 g for 10
100 minutes and the plasma/serum snap frozen in liquid nitrogen and stored at -80°C prior to batch
101 analysis.

102

103 ***Lipids, Glucose and Insulin***

104 Triglycerides and glucose were assessed using photometry (Randox Daytona Plus, Randox,
105 County Antrim, UK). Insulin was assessed via radio-immunoassay (Invitron Insulin Assay Kit
106 (IV2-001), Invitron Ltd, UK). The intra and inter-assay coefficient of variations (CVs) were
107 both < 5% for all measures.

108

109 ***Nitric Oxide Bioavailability***

110 Plasma nitrite (NO_2^-) was determined using ozone-based chemiluminescence (OBC Model
111 280i, NOA, Sievers, Boulder, CO, USA) following reduction by sodium iodide in acetic acid.
112 The intra and inter-assay CVs were both < 5%.

113

114 ***Oxidative Stress***

115 The ascorbate free radical ($\text{A}^{\cdot-}$) was employed as a direct measure of global oxidative stress as
116 previously described (11). Plasma was injected into a high-sensitivity multiple-bore sample
117 cell (AquaX; Bruker Daltonics Inc., Billerica, MA, USA) housed within an electron
118 paramagnetic resonance (EPR) spectrometer operating at X-band (9.87 GHz). The intra and
119 inter-assay CVs were both < 5%.

120

121

122 *Statistical Analysis*

123 Following confirmation of distribution normality (Shapiro-*W*-Wilk tests), data were analysed
124 using a 2-way repeated measures (pre-meal *vs.* post-meal and young *vs.* old) analysis of
125 variance. Where interaction effects were identified, differences were located using *Bonferonni*-
126 corrected paired samples *t*-tests and independent samples *t*-tests. Significance was established
127 at $P < 0.05$ and data expressed as mean \pm standard deviation.

128

129 Results

130 At baseline, the aged presented with lower MCAv and higher fasted triglycerides, glucose,
131 insulin, systolic blood pressure, pulse pressure, CVR_{CO_2Hyper} and corresponding CVR_{CO_2} range
132 ($P < 0.05$ vs. young, Table and Figure). During PPH, circulating triglycerides, glucose and
133 insulin increased in both the young (triglycerides: 0.85 ± 0.50 to 2.22 ± 1.51 mmol/L; glucose:
134 5.48 ± 0.63 to 5.80 ± 0.63 mmol/L; insulin: 39 ± 19 to 97 ± 64 pmol/L, all $P < 0.05$ vs. pre-
135 meal) and aged (triglycerides: 1.47 ± 0.70 to 3.02 ± 1.62 mmol/L; glucose: 6.07 ± 0.51 to 6.70
136 ± 1.03 mmol/L; insulin: 54 ± 30 to 148 ± 91 pmol/L; all $P < 0.05$ vs. pre-meal, Figure Panels
137 A, B, C). This was associated with a selective increase in A^+ in the young but not the aged (P
138 < 0.05 vs. pre-meal, Figure Panel D), whereas NO_2^- was shown to decrease in both groups (P
139 < 0.05 vs. pre-meal, Figure Panel E). PPH had no effect on systolic blood pressure, diastolic
140 blood pressure, MAP or pulse pressure ($P > 0.05$ vs. pre-meal, Table), but was associated with
141 a selective reduction in CVR_{CO_2Hyper} and corresponding CVR_{CO_2} range in the aged ($P < 0.05$
142 vs. pre-meal, Figure Panel F and Table).

143 Discussion

144 The major finding of the present study is that PPH was shown to selectively impair CVR_{CO_2} in
145 the aged, but not in the young adults and this was associated with a systemic reduction in
146 bioactive NO_2^- in the absence of any systemic increase in free radical formation. This has
147 important clinical implications given that an increasing proportion of the world's population is
148 consuming high-fat diets and thus will spend large proportions of the day in a PPH state.

149 To the best of our knowledge, this is the first study to investigate the effect of PPH on the
150 human cerebrovasculature across the adult lifespan. Our findings are in agreement with those
151 by Dunn and Walter (9) who reported that PPH had no effect on the brain in young adults. Our
152 study has extended these findings demonstrating for the first time that the aged brain is more
153 vulnerable to the adverse effects of PPH in the form of reduced CVR_{CO_2} , independent of any
154 baseline differences in BMI between age groups. Thus, it would appear that the acute effects
155 of PPH are not simply confined to the systemic circulation (2-5), but can equally impact the
156 local cerebral circulation. Interestingly, baseline CVR_{CO_2} was higher in the aged compared to
157 the young, which is in agreement with recent reports (12). Nonetheless, the reductions in
158 CVR_{CO_2} observed post-prandially in the aged, combined with the global reduction in MCA_v ,
159 may enhance stroke risk (8) and increase vulnerability to neurodegenerative diseases (13). It is
160 therefore important that nutritional guidelines are adhered to, which promote a varied and
161 nutrient dense diet that is low in fat (14).

162 We sought to provide a mechanistic basis to explain these findings through specific assessment
163 of systemic oxidative-nitrosative stress given that vascular endothelial dysfunction at least in
164 the systemic and pulmonary circulation, has traditionally been attributed to a free radical-
165 mediated reduction in endothelium-derived vascular NO bioavailability (11, 15). The observed
166 changes in plasma NO_2^- support the concept that reductions in cerebrovascular function
167 following a high-fat meal were mediated by a reduction in NO bioavailability, at least in the

168 aged. However, this does not appear to be caused by an increase in oxidative stress, since no
169 additional elevations in A^+ were observed post-prandially. It was interesting to note that the
170 baseline A^+ to which the aged were chronically exposed (*i.e.* prior to ingestion of the high-fat
171 meal) was identical in magnitude to the acute increase observed post-prandially in the young,
172 placing the degree of oxidative stress into clear physiological context, perhaps reflecting an
173 “upper-limit” of oxidation. Alternatively, the net loss of plasma NO_2^- from the peripheral
174 circulation may equally reflect increased consumption and concomitant liberation of NO or
175 dinitrogen trioxide catalysed by acidic disproportionation (16) in an attempt to “salvage”
176 vascular endothelial function in the chronically oxidatively-stressed cerebrovasculature of the
177 older adults to maintain function.

178 Similar impairments in vascular function independent of changes in oxidative stress, albeit
179 using indirect biomarkers of free radical-mediated lipid peroxidation have previously been
180 reported (17, 18), though the underlying mechanisms remain to be established. A possible
181 explanation may relate to the length of time that the aged brain has been exposed to increased
182 levels of oxidative-stress across the lifespan (*i.e.* much longer than the young). This “chronic”
183 effect may subsequently “blunt” the sensitivity of the systemic and cerebral vasculature to NO,
184 a concept that has not previously been interrogated in this setting to the best of our knowledge.
185 Thus, when the aged are challenged with the “additional” oxidative-nitrosative stress imposed
186 by the high-fat meal, the vasculature is unable to synthesise sufficient NO and/or respond
187 sufficiently to maintain function, resulting in a decrease in CVR_{CO_2} . What is apparent is that
188 the reduction in NO_2^- and subsequent impairment in vascular function observed in the aged are
189 unlikely to be explained by changes in vascular compliance, since there were no changes in
190 systolic blood pressure, diastolic blood pressure or pulse pressure following consumption of
191 the high-fat meal. Similarly, they are unlikely to be explained by any differences in the
192 metabolic responses to the meal, since the relative increase in triglycerides, glucose and insulin

193 were comparable between both young and aged. Though, it is likely that the slightly elevated
194 levels of triglycerides and glucose observed at baseline in the aged contributed to their chronic
195 levels of oxidative stress, given the additional free radical production associated with their
196 metabolism in larger quantities (19). These findings further highlight that additional research
197 is warranted to better understand the causes of the deterioration in vascular function following
198 the consumption of a high-fat meal, particularly in the aged.

199 A potential limitation to our study relates to between group differences in baseline measures,
200 an almost unavoidable consequence when recruiting participants at polar opposites of the
201 ageing continuum. However, the primary focus of the study was to determine within-group
202 responses to the high-fat meal, rather than focus on the between-group responses. Thus,
203 reducing the importance of having to match groups for anthropometrics, as well metabolic and
204 vascular function. Our study would have also benefited by including an additional measure of
205 general vascular reactivity such as flow-mediated dilatation to help explain our findings, since
206 it is in part NO-dependant (20), similar to CVR_{CO_2} (7). However, given the established
207 literature that demonstrates a reduction in general vasoreactivity following consumption of a
208 high-fat meal (2-5), it is plausible to assume that similar changes occurred in the present study.

209 In conclusion, the present findings demonstrate that a single high-fat meal can impair
210 cerebrovascular function in the old but not in the young mediated in part by a systemic
211 reduction in vascular NO bioavailability that was independent of any “additional” elevation in
212 oxidative stress or changes in vascular compliance. These findings have important implications
213 given the established relationship between impaired CVR_{CO_2} and increased stroke risk.

214 Acknowledgements

215 We appreciate the commitment of the participants. Technical support was provided by Mr.
216 Gareth Dunseath.

217

218 Declarations of Interest

219 The authors declare that there are no competing interests associated with the manuscript.

220

221 Funding

222 Damian Bailey is a Royal Society Wolfson Research Fellow (#WM170007). Damian Bailey
223 received a Fellowship Award for Christopher Marley from the JPR Williams Trust.

224

225 Author Contributions

226 Damian Bailey and Christopher Marley led the acquisition/analysis/interpretation of data and
227 wrote the original paper. Danielle Hodson and Julien Brugniaux made substantial
228 contributions to data acquisition and gave final approval of the version to be submitted. Lewis
229 Fall made substantial contributions to data analysis and gave final approval of the version to
230 be submitted. Damian Bailey was the Principal Investigator, recipient of funding and critically
231 revised the original paper for important intellectual content and gave final approval of the
232 version to be submitted including any revisions thereof.

233

234 Clinical Perspectives

- 235 i. Post-prandial hyperlipidaemia has been shown to acutely impair systemic vascular
236 endothelial function, potentially attributable to increased oxidative-nitrosative stress. It
237 remains to be determined whether this extends to the cerebrovasculature.
- 238 ii. Our findings identified that post-prandial hyperlipidaemia impairs cerebrovascular function
239 in the aged, but not in the young and this was associated with a systemic reduction in nitric
240 oxide bioavailability in the absence of increased oxidative stress or changes in vascular
241 compliance.
- 242 iii. These findings have important clinical consequences since an increasing proportion of the
243 world's population is consuming high-fat diets and an impairment in cerebrovascular
244 function is associated with increased stroke risk and vulnerability to neurodegenerative
245 diseases.

246 References

- 247 1. Patsch JR, Karlin JB, Scott LW, Smith LC, Gotto AM, Jr. Inverse relationship between
248 blood levels of high density lipoprotein subfraction 2 and magnitude of postprandial lipemia.
249 Proceedings of the National Academy of Sciences of the United States of America.
250 1983;80(5):1449-53.
- 251 2. Marchesi S, Lupattelli G, Schillaci G, Pirro M, Siepi D, Roscini AR, et al. Impaired
252 flow-mediated vasoactivity during post-prandial phase in young healthy men. *Atherosclerosis*.
253 2000;153(2):397-402.
- 254 3. Tsai WC, Li YH, Lin CC, Chao TH, Chen JH. Effects of oxidative stress on endothelial
255 function after a high-fat meal. *Clin Sci (Lond)*. 2004;106(3):315-9.
- 256 4. Vogel RA, Corretti MC, Plotnick GD. Effect of a single high-fat meal on endothelial
257 function in healthy subjects. *The American journal of cardiology*. 1997;79(3):350-4.
- 258 5. West SG, Hecker KD, Mustad VA, Nicholson S, Schoemer SL, Wagner P, et al. Acute
259 effects of monounsaturated fatty acids with and without omega-3 fatty acids on vascular
260 reactivity in individuals with type 2 diabetes. *Diabetologia*. 2005;48(1):113-22.
- 261 6. Bailey DM, Evans KA, McEneny J, Young IS, Hullin DA, James PE, et al. Exercise-
262 induced oxidative-nitrosative stress is associated with impaired dynamic cerebral
263 autoregulation and blood-brain barrier leakage. *Experimental physiology*. 2011;96(11):1196-
264 207.
- 265 7. Lavi S, Gaitini D, Milloul V, Jacob G. Impaired cerebral CO₂ vasoreactivity:
266 association with endothelial dysfunction. *American journal of physiology Heart and circulatory*
267 *physiology*. 2006;291(4):H1856-61.
- 268 8. Gupta A, Chazen JL, Hartman M, Delgado D, Anumula N, Shao H, et al.
269 Cerebrovascular reserve and stroke risk in patients with carotid stenosis or occlusion: a

- 270 systematic review and meta-analysis. *Stroke; a journal of cerebral circulation.*
271 2012;43(11):2884-91.
- 272 9. Dunn WG, Walters MR. A randomised crossover trial of the acute effects of a deep-
273 fried Mars bar or porridge on the cerebral vasculature. *Scott Med J.* 2014;59(4):182-7.
- 274 10. Bailey DM, Jones DW, Sinnott A, Brugniaux JV, New KJ, Hodson D, et al. Impaired
275 cerebral haemodynamic function associated with chronic traumatic brain injury in professional
276 boxers. *Clin Sci (Lond).* 2013;124(3):177-89.
- 277 11. Bailey DM, Dehnert C, Luks AM, Menold E, Castell C, Schendler G, et al. High-
278 altitude pulmonary hypertension is associated with a free radical-mediated reduction in
279 pulmonary nitric oxide bioavailability. *The Journal of physiology.* 2010;588(Pt 23):4837-47.
- 280 12. Zhu YS, Tarumi T, Tseng BY, Palmer DM, Levine BD, Zhang R. Cerebral vasomotor
281 reactivity during hypo- and hypercapnia in sedentary elderly and Masters athletes. *Journal of*
282 *cerebral blood flow and metabolism : official journal of the International Society of Cerebral*
283 *Blood Flow and Metabolism.* 2013;33(8):1190-6.
- 284 13. Gorelick PB, Scuteri A, Black SE, Decarli C, Greenberg SM, Iadecola C, et al. Vascular
285 contributions to cognitive impairment and dementia: a statement for healthcare professionals
286 from the american heart association/american stroke association. *Stroke; a journal of cerebral*
287 *circulation.* 2011;42(9):2672-713.
- 288 14. Public Health England. A Quick Guide to the Government's Healthy Eating
289 Recommendations 2017 [Available from:
290 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/595133/A_qui](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/595133/A_quick_guide_to_govt_healthy_eating.pdf)
291 [ck_guide_to_govt_healthy_eating.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/595133/A_quick_guide_to_govt_healthy_eating.pdf).
- 292 15. Gryglewski RJ, Palmer RM, Moncada S. Superoxide anion is involved in the
293 breakdown of endothelium-derived vascular relaxing factor. *Nature.* 1986;320(6061):454-6.

- 294 16. Bailey DM, Rasmussen P, Overgaard M, Evans KA, Bohm AM, Seifert T, et al. Nitrite
295 and S-Nitrosohemoglobin Exchange Across the Human Cerebral and Femoral Circulation:
296 Relationship to Basal and Exercise Blood Flow Responses to Hypoxia. *Circulation*.
297 2017;135(2):166-76.
- 298 17. Gopaul NK, Zacharowski K, Halliwell B, Anggard EE. Evaluation of the postprandial
299 effects of a fast-food meal on human plasma F(2)-isoprostane levels. *Free Radic Biol Med*.
300 2000;28(5):806-14.
- 301 18. Williams MJ, Sutherland WH, McCormick MP, de Jong SA, Walker RJ, Wilkins GT.
302 Impaired endothelial function following a meal rich in used cooking fat. *Journal of the*
303 *American College of Cardiology*. 1999;33(4):1050-5.
- 304 19. Roberts CK, Vaziri ND, Wang XQ, Barnard RJ. Enhanced NO inactivation and
305 hypertension induced by a high-fat, refined-carbohydrate diet. *Hypertension*. 2000;36(3):423-
306 9.
- 307 20. Green DJ, Dawson EA, Groenewoud HM, Jones H, Thijssen DH. Is flow-mediated
308 dilation nitric oxide mediated?: A meta-analysis. *Hypertension*. 2014;63(2):376-82.
309

Table. Changes in cerebrovascular function following a high-fat meal.

Age Group	Young (<i>n</i> = 19)		Aged (<i>n</i> = 19)		<i>P</i> Values		
	Pre-Meal	Post-Meal	Pre-Meal	Post-Meal	Age	Meal	Age × Meal
MCAv (cm.s⁻¹)	62 ± 13	61 ± 10	48 ± 12	46 ± 10	0.00	0.37	0.67
Systolic BP (mmHg)	123 ± 10	122 ± 9	132 ± 16	129 ± 16	0.04	0.46	0.50
Diastolic BP (mmHg)	70 ± 10	68 ± 8	70 ± 11	68 ± 10	0.98	0.24	0.84
MAP (mmHg)	87 ± 9	86 ± 7	91 ± 11	89 ± 10	0.85	0.40	0.72
Pulse Pressure (mmHg)	52 ± 7	54 ± 9	62 ± 14	61 ± 13	0.02	0.69	0.28
CVR_{CO2Hyper} (%.mmHg⁻¹)	2.68 ± 0.85	2.39 ± 0.85	5.46 ± 1.71†	4.07 ± 1.06*†	0.00	0.00	0.00
CVR_{CO2Hypo} (%.mmHg⁻¹)	2.49 ± 0.45	2.12 ± 0.58	2.47 ± 0.89	2.28 ± 1.29	0.77	0.05	0.51
CVR_{CO2} range (%.mmHg⁻¹)	5.17 ± 0.90	4.51 ± 1.16	8.21 ± 1.95†	6.35 ± 1.61*†	0.00	0.00	0.01

Values are mean ± SD; MCAv, middle cerebral artery velocity; BP, blood pressure; MAP, mean arterial blood pressure;

CVR_{CO2Hyper}/CVR_{CO2Hypo}, cerebrovascular reactivity to hypercapnea/hypocapnea; CVR_{CO2} range, combination of CVR_{CO2Hyper} and

CVR_{CO2Hypo}. */† = difference within/between age groups (*P* < 0.05)




Figure. Metabolic (A-E) and haemodynamic (F) responses to a high-fat meal.

Data are mean \pm SD; Young $n = 19$, Aged $n = 19$; *different vs. pre-meal for given Group ($P < 0.05$), †different vs. Group ($P < 0.05$).

24

25

26