



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

© 2018, *Appetite*. This is an author produced version of a paper accepted for publication in *Appetite*. Uploaded in accordance with the publisher's self- archiving policy.

Buckland, N., & Dalton, M. (2018). Commentary: Methodological and reporting practices for laboratory studies assessing food intake using fixed and ad libitum test meals. *Appetite*. XX(X).

UNCORRECTED PROOFS

22

23

24 **Commentary: Methodological and reporting practices for laboratory studies assessing food intake**
25 **using fixed and ad libitum test meals.**

26 Nicola J. Buckland, N.J.^{1*} & Michelle Dalton²

27 ^{1*}Department of Psychology, University of Sheffield, Cathedral Court, 1 Vicar Lane, Sheffield, S1 2LT

28 ²School of Social and Health Sciences, Leeds Trinity University, Brownberrie Lane, Leeds, LS18 5HD

29 Keywords: Laboratory food intake, methods; satiation, satiety, standardised test meal

30 *Corresponding author: Dr Nicola Buckland, email: n.buckland@sheffield.ac.uk, telephone: (+44) 114

31 222 6508

32

UNCORRECTED PROOFS

33 A number of expert reports have provided methodological recommendations on how to conduct
34 rigorous and scientifically sound laboratory studies to investigate appetite control (most recently:
35 Blundell et al., 2010, Blundell et al., 2009, Gibbons et al., 2014). However, a recent examination of the
36 methodologies used in laboratory food intake studies by Robinson, Bevelander, Field, and Jones
37 (2018) showed that many failed to adopt basic methodological procedures and reporting practices.
38 Based on their examination Robinson et al. proposed recommendations that should be adopted as
39 best practice in appetite-related research. We wholly support Robinson et al.'s effort to highlight
40 where scientific rigour needs to be improved in this research area. Indeed, in one of our recent meta-
41 analyses on studies assessing food intake we also concluded that most studies were of low
42 methodological quality (Buckland et al., *under review*).

43 While we support the recommendations of Robinson et al. (2018), we propose that in their current
44 form the recommendations are limited and overlook other important '*basic*' methodological factors
45 that should be considered when designing and reporting studies that assess food intake. Such factors
46 include adopting additional pre-manipulation control procedures (e.g. controlling for alcohol intake
47 and physical activity levels), designing (and reporting) an appropriate order for study procedures to
48 ensure that any manipulations are not confounded by other study measures (e.g. weighing
49 participants or administering psychometric questionnaires before assessing food intake), detailed
50 reporting of sample type (e.g. student, community-based, dieting status), reporting whether measures
51 taken were objective or subjective (e.g. self-reported versus researcher measured body weight and
52 height) and appropriate design and reporting of standardised test meals. Reporting information on
53 these additional methodological factors would facilitate the replication of studies. Further,
54 recommendations may need to be tailored according to study aims. For example, when considering
55 the issue of standardised test meals, recommendations will vary if the study is examining processes
56 affecting satiation or satiety or if it is examining food hedonics and food choice.

57 A thorough review and examination of each of these additional factors are beyond the scope of this
58 commentary, so we will focus on extending Robinson et al. (2018) recommendations with regards to
59 developing criteria for appropriate standardised test meal design.

60 **Appropriate design and reporting of standardised test meals**

61 The focus of Robinson et al. (2018) examination was on laboratory studies of human food intake. The
62 advantage of laboratory assessments of eating behaviour is that they allow for the precise assessment
63 of food intake in a controlled environment that is free from potential confounding variables such as
64 extraneous smells, sounds, competing activities and social stimuli (Blundell et al., 2009). Broadly
65 speaking there are two forms of food intake assessment within the laboratory; the first is a measure
66 of fixed intake (termed as “fixed energy meals”) where the type and amount of food consumed by the
67 participant is pre-determined by the researcher and is less susceptible to confounding variables. The
68 second is a measure of ad libitum intake where the amount (and in some cases type) of food consumed
69 is determined by the participant (ideally in response to the experimental manipulation) within the
70 limitations of the experimental design. This second measure is more vulnerable to confounding factors
71 (Stubbs et al., 1997). For both fixed and ad libitum test meals the type and amount of food provided
72 requires careful consideration as variation in these factors has been shown to influence the amount
73 of food consumed (Beaulieu et al., 2017; Hetherington & Blundell-Birtill, 2018). Within their
74 examination, Robinson et al. assessed whether the studies reported the types of foods provided but
75 they did not provide recommendations on which variables are important to consider when designing
76 and reporting fixed and ad libitum test meals used in laboratory studies of human food intake.

77 **Fixed energy meals**

78 Fixed energy test meals are those in which the researcher provides the participant with a compulsory
79 “fixed” test meal that they are instructed to consume in its entirety. Fixed energy test meals allow for
80 the composition of food to be manipulated and standardised across participants. Fixed energy test
81 meals allow for increased experimental control in designs where food is being used as an independent

82 variable. However, fixed energy test meals are not suitable for studies examining satiation as they do
83 not account for individual differences in energy requirements.

84 Fixed energy test meals are also useful to standardise participants' appetite before they are exposed
85 to an experimental manipulation. When used to standardise appetite, ideally fixed meals should be
86 tailored to individual daily energy needs (e.g. based on Schofield equations or measured resting
87 metabolic rate). The proportion of daily energy requirements a fixed meal provides will be determined
88 by study aims and time of day the test meal is served (Dalton et al., 2015). An alternative method
89 when there are multiple conditions is to have participants self-determine their fixed meal by providing
90 an ad libitum amount in the first condition and asking them to eat to comfortable fullness. The amount
91 consumed can then be provided in the experimental conditions that follow (for an example see
92 Beaulieu et al., 2017). It is important to consider individual energy requirements as providing the same
93 portion to all participants does not account for energy needs differing depending on individual
94 characteristics such as age, gender, body weight and body composition (Ravussin & Bogardus, 1989).
95 This may lead to some participants receiving too little and still feeling hungry and others receiving too
96 much and feeling too full which can interfere with any subsequent assessments of food intake.

97 **Ad libitum test meals**

98 In ad libitum test meal designs participants are normally provided with a larger than can be consumed
99 portion of food, which the researcher weighs before and after consumption. A range of foods are
100 often provided for participants to choose from which allows for the assessment of quantitative aspects
101 of eating behavior (i.e. how much) and qualitative aspects of eating behavior (i.e. nutrient and/or
102 sensory food choice). When used correctly ad libitum test meals are useful to assess the process of
103 satiation (i.e. meal size and termination) however there are several important considerations when
104 designing ad libitum test meals. Research has shown that factors such as variety, texture, physical
105 form (liquid or solid), palatability and energy density can induce over- and under-eating in laboratory
106 conditions (Buckland et al., *in press*; de Graaf, 2012; Hetherington, Foster, Newman, Anderson &

107 Norton, 2006; Raynor & Epstein, 2001; Rolls, Van Duijvenvoorde, & Rolls, 1984). Additionally, care
108 must be taken with regards to the portion size of the ad libitum test meal items as larger portion sizes
109 have been shown to lead to greater intake (for a review see Hetherington & Blundell-Birtill, 2018;
110 Zlatevska et al., 2014) whereas providing small portions may constrain participants' food intake and
111 limit the opportunity to observe effects of the independent variable. It is recommended that the
112 portion size of ad libitum test meals is clearly reported in each study and the range of food consumed
113 is provided.

114 Furthermore, it is recommended that researchers assess participants' liking of study foods as an
115 inclusion criterion; a factor that was not examined by Robinson et al. (2018). Liking for food has a
116 positive effect on food intake (De Graaf et al., 1999) and therefore to accurately assess the effects of
117 a manipulation, the foods provided must be liked by participants (Blundell et al., 2010). If study foods
118 differ between study conditions, then food liking should be matched across conditions to ensure any
119 differences in intake can be attributed to the study manipulation rather than the extent to which
120 participants like the food. For example, one study compared whether intake differed if participants
121 were provided with the same (fish and chips or beef stew) or different (lemon mousse) food to that
122 previously eaten (Ferriday et al., 2016). Compared to when eating the same food, participants ate less
123 of the different food and reported feeling less full. Crucially, the authors did not check pre-study
124 whether participants liked the lemon mousse (any pre-screening attempts were not reported) and as
125 such as the authors discussed, it was unclear whether participants ate less of the mousse because of
126 the study manipulation (varied the test foods to be either the same or different to foods previously
127 eaten) or due to a dislike for the lemon mousse. Such issues can easily be prevented by assessing liking
128 for study foods in a pre-study screening questionnaire, with the aim of including low liking for the
129 study foods (e.g. ratings of <4 on a 7-point Likert scale) as an exclusion criterion (Gibbons et al., 2014)
130 In addition, study foods should adequately undergo pilot testing to ensure they are equally palatable.

131 Lastly, studies should also report the test meal environment, including the presence or absence of
132 social others, participants' focus on the test meal and the time of day that test meals were
133 administered. The presence of social others (social influences) has been shown to influence food
134 intake (Herman et al., 2003). As such, if social cues are not part of the research question then
135 participants should be tested in individual cubicles. Distractions such as watching television, listening
136 to audiobooks and completing computer tasks increase food intake (Oldham-Cooper et al., 2011, Higgs
137 and Woodward, 2009, Bellisle et al., 2004). Therefore, food intake should also be assessed in a
138 distraction-free environment where participants do not have access to their mobile phone, computer
139 or other distractions to ensure their attention is focussed on the test meal. The time of day that the
140 test session takes place can also influence food intake. Certain foods will be more culturally
141 appropriate at particular times of the day compared to others. As such, to avoid confounding the
142 variable of interest, test foods should be appropriate for the time of day that the test session takes
143 place (Blundell et al., 2010).

144 These methodological aspects related to study foods are not exhaustive of the "basic" methods that
145 researchers should consider when assessing food intake within the laboratory using standardized test
146 meals (see Blundell et al., 2010). We have raised these points to demonstrate that Robinson et al.
147 (2018) did not discuss or provide recommendations for a large number of "basic" methodological and
148 reporting practices. While we are aware that Robinson et al. acknowledged that "*it was not feasible*
149 *(however), to evaluate all aspects of study design and reporting*" (p.490) we believe that providing
150 restricted recommendations risks future studies overlooking important methods. Overlooking such
151 methods can lead to the collection of low quality data and make it difficult to form justifiable
152 conclusions (Brown et al., 2018). As such, in line with Robinson, we call for experts in the laboratory
153 assessment of food intake to agree and establish a comprehensive set of recommendations that can
154 be used by researchers and reviewers of manuscripts to encourage and promote scientifically sound
155 research.

156 Conclusions

157 We support Robinson et al. (2018) recommendations to promote scientific rigour in laboratory studies
158 investigating food. However, to avoid important aspects of research design being overlooked we
159 strongly urge experts in eating behaviour to collaboratively establish more thorough
160 recommendations.

161 **References**

- 162 BEAULIEU, K., HOPKINS, M., BLUNDELL, J. & FINLAYSON, G. 2017 Impact of physical activity and
163 dietary fat content on passive overconsumption of energy in non-obese adults. *International Journal*
164 *of Behavioral Nutrition and Physical Activity*, 14:14
- 165 BELLISLE, F., DALIX, A. M. & SLAMA, G. 2004. Non food-related environmental stimuli induce
166 increased meal intake in healthy women: comparison of television viewing versus listening to a
167 recorded story in laboratory settings. *Appetite*, 43, 175-180.
- 168 BLUNDELL, J., DE GRAAF, C., HULSHOF, T., JEBB, S., LIVINGSTONE, B., LLUCH, A., MELA, D., SALAH, S.,
169 SCHURING, E., VAN DER KNAAP, H. & WESTERTERP, M. 2010. Appetite control: methodological
170 aspects of the evaluation of foods. *Obesity Reviews*, 11, 251-270.
- 171 BLUNDELL, J. E., FINLAYSON, G., HALFORD, J., HETHERINGTON, M. M. & KING, N. (eds.) 2009.
172 *Measuring food intake, hunger, satiety and satiation in the laboratory*, London: Sage.
- 173 BROWN, A. W., KAISER, K. A. & ALLISON, D. B. 2018. Issues with data and analyses: Errors, underlying
174 themes, and potential solutions. *Proceedings of the National Academy of Sciences of the United*
175 *States of America*, 115, 2563-2570.
- 176 BUCKLAND, N. J., ER, V., REDPATH, I. & BEAULIEU, K. *under review*. Priming food intake with weight
177 control cues: A systematic review with a meta-analysis. *International Journal of Behavioral Nutrition*
178 *and Physical Activity*.
- 179 DALTON, M., HOLLINGWORTH, S., BLUNDELL, J. & FINLAYSON, G. Weak satiety responsiveness is a
180 reliable trait associated with hedonic risk factors for overeating among women. *Nutrients*, 7(9),
181 7421-7436
- 182 DE GRAAF, C., DE JONG, L. S. & LAMBERS, A. C. 1999. Palatability affects satiation but not satiety.
183 *Physiology & Behavior*, 66, 681-688.
- 184 DE GRAAF, C. 2012. Texture and satiation: The role of oro-sensory exposure time. *Physiology &*
185 *Behavior*, 107, 496-501.
- 186 FERRIDAY, D., BOSWORTH, M. L., GODINOT, N., MARTIN, N., FORDE, C. G., VAN DEN HEUVEL, E.,
187 APPLETON, S. L., MOSS, F. J. M., ROGERS, P. J. & BRUNSTROM, J. M. 2016. Variation in the Oral
188 Processing of Everyday Meals Is Associated with Fullness and Meal Size; A Potential Nudge to Reduce
189 Energy Intake? *Nutrients*, 8.
- 190 GIBBONS, C., FINLAYSON, G., DALTON, M., CAUDWELL, P. & BLUNDELL, J. E. 2014. Studying eating
191 behaviour in humans. *Journal of Endocrinology*, 222, G1-G12.

- 192 HERMAN, C. P., ROTH, D. A. & POLIVY, J. 2003. Effects of the presence of others on food intake: A
193 normative interpretation. *Psychological Bulletin*, 129, 873-886.
- 194 HETHERINGTON, M.M., FOSTER, R., NEWMAN, T., ANDERSON, A & NORTON, G. 2006 Understanding
195 variety: Tasting different foods delays satiation. 87(2), 263-271
- 196 HETHERINGTON, M.M. & BLUNDELL-BIRTILL (2018) The portion size effect and overconsumption
197 towards downsizing solutions for children and adolescents. *Nutrition Bulletin*, 43, 61-68
- 198 HIGGS, S. & WOODWARD, M. 2009. Television watching during lunch increases afternoon snack
199 intake of young women. *Appetite*, 52, 39-43.
- 200 OLDHAM-COOPER, R. E., HARDMAN, C. A., NICOLL, C. E., ROGERS, P. J. & BRUNSTROM, J. M. 2011.
201 Playing a computer game during lunch affects fullness, memory for lunch, and later snack intake.
202 *American Journal of Clinical Nutrition*, 93, 308-313.
- 203 RAVUSSIN, E. & BOGARDUS, C. 1989. Relationships of genetics, age, and physical activity fitness to
204 daily energy expenditure and fuel utilisation. *American Journal of Clinical Nutrition*. 5(1), 968-975
- 205 RAYNOR, H.A. & EPSTEIN, L.H. 2001 Dietary variety, energy regulation and obesity. *Psychological*
206 *Bulletin*. 127(3), 325-341
- 207 ROLLS, B.J., VAN DUIJVENVOORDE, P.M. & ROLLS, E.T. 1984 Pleasantness changes and food intake in
208 a varied four course meal. *Appetite*, 5(4), 337-348
- 209 ROBINSON, E., BEVELANDER, K. E., FIELD, M. & JONES, A. 2018. Methodological and reporting quality
210 in laboratory studies of human eating behavior. *Appetite*.
- 211 STUBBS, R.J., JOHNSTONE, A.M., O'REILLY, L.M. & POPPITT, S.D. 1997 Methodological issues relating
212 to the measurement of food, energy and nutrient intake in human laboratory-based studies.
213 *Proceedings of the Nutrition Society*. 57(3), 357-372
- 214 YEOMANS, M. R. 2018. Measuring Appetite and Food Intake *Methods in Consumer Research, Volume*
215 *2* (pp. 119-149): Elsevier.
- 216 ZLATEVSKA, N., DUBELAAR, C. & HOLDEN, S.S. 2014 Sizing up the effect of portion size on
217 consumption: a meta-analytic review. *Journal of Marketing*, 78(3), 140-154