



© 2019 Taylor & Francis. This is an Accepted Manuscript of an article published by Taylor & Francis in The Journal of Sports Sciences on 3<sup>rd</sup> June 2019 available online:

<http://www.tandfonline.com/10.1080/02640414.2019.162561>

3

Birdsey, L.P. et al. (2019). Neuromuscular, physiological and perceptual responses to an elite netball tournament. *Journal of Sports Sciences*. 37(19), pp. 2169-2174

1 Full title: Neuromuscular, physiological and perceptual responses to an elite  
2 netball tournament.

3

4 Running title: Neuromuscular, physiological and perceptual responses to elite  
5 netball

6

7 Laurence P. Birdsey<sup>ab</sup>, Matthew Weston<sup>c</sup>, Mark Russell<sup>d</sup>, Michael Johnston<sup>e</sup>,  
8 Christian J. Cook<sup>f</sup> and Liam P.Kilduff<sup>a\*</sup>

9 *<sup>a</sup>A-STEM College of Engineering, Swansea University, Swansea, UK; <sup>b</sup>Sport*

10 *Wales, Cardiff, UK; <sup>c</sup>Department of business and law, Teesside University,*

11 *Middlesborough, UK; <sup>d</sup>School of social and health sciences, Leeds trinity*

12 *University, UK; <sup>e</sup>British Athletics, Birmingham, UK; <sup>f</sup>School of sport, health and*

13 *exercise sciences, Bangor University, UK*

14 Corresponding author: Liam P. Kilduff, A-STEM College of engineering,

15 Swansea University Bay Campus, Fabian Way, Crymlyn Burrows, Swansea,

16 SA81 8EN. L.kilduff@swansea.ac.uk

17

18 Word count: 3354

19

20

21

22

23

24 **Neuromuscular, physiological and perceptual responses to an elite**  
25 **netball tournament**

26 **Abstract**

27 To examine responses to an International netball tournament, female athletes  
28 ( $n=11$ ) played three matches over consecutive days. External (accelerometry) and  
29 internal (heart rate; HR, session; sRPE, and differential; dRPE, rating of perceived  
30 exertion) load measures quantified match intensity. On match-day mornings, and  
31 three days after match three, well-being (brief assessment of mood; BAM+),  
32 biochemical (creatinine kinase concentration; CK), neuromuscular (jump height; JH,  
33 peak power output; PPO) and endocrine function (salivary cortisol; C, testosterone;  
34 T, concentrations) were assessed. External load was similar between matches  
35 whereas dRPE and sRPE was greatest for match three. Following match one, CK  
36 increased, whereas BAM+, JH, C and T decreased. Following two matches, BAM+,  
37 PPO, and T decreased with CK increasing versus baseline. Following consecutive  
38 matches, CK (likely moderate;  $27.9\% \pm 19.5\%$ ) and C (possibly moderate;  $43.3\%$   
39  $\pm 46.8\%$ ) increased, whilst BAM+ (possibly moderate;  $-20.6\% \pm 24.4\%$ ) decreased.  
40 Three days post-tournament BAM+, T, PPO, and JH decreased. Mid-court elicited  
41 higher mean HR (possibly moderate;  $3.7\% \pm 3.8\%$ ), internal and external intensities  
42 (possibly very large;  $85.7\% \pm 49.6\%$ ) compared with goal-based positions.  
43 Consecutive matches revealed a dose-response relationship for well-being and  
44 physiological function; a response evident three days post-tournament.

45 **Keywords:** recovery; monitoring; load; team sport; readiness to train

46 **Introduction**

47 Whilst several studies have reported the movement demands of elite netball in  
48 recent years (Bailey, Gustin, Mackey, & Dwyer, 2017; Fox, Spittle, Otago, &  
49 Saunders, 2013; Young, Gustin, Sanders, Mackey, & Dwyer, 2016), to date no  
50 studies have profiled the physiological responses to elite level tournament match-  
51 play. Indeed, only three studies have reported the movement demands of elite  
52 netball, one by use of notational analysis (Fox et al., 2013) and two by  
53 accelerometry (Bailey et al., 2017; Young et al., 2016). Goal defence (GD),  
54 goalkeeper (GK) and goal shooter (GS) positions were reported to perform at the  
55 the lowest playing intensities and highest proportions of match time spent in the  
56 low-intensity zones when compared to players occupying wing attack (WA), wing  
57 defence (WD), centre (C), and goal attack (GA) positions (Young et al., 2016).  
58 Additionally, Bailey et al., (2017) reported the accelerometer-based loads  
59 associated with typical activities, reporting off-ball guarding to elicit the highest  
60 load per instance, whilst jogging accumulated the greatest load across a match.

61

62 At present, a single study has reported the responses to an isolated match reporting  
63 a reduction in perception of fatigue and neuromuscular performance immediately  
64 and 24 h after an 80 min elite level match, returning to baseline 36 h later (Wood,  
65 Kelly, & Gabbett, 2013). Many tournaments require teams to play up to eight  
66 matches in 10 days, therefore, the demands are not limited to that of a single match,  
67 rather the ability to perform and recover over a series of days. Findings of previous  
68 studies reporting the neuromuscular and perceptual recovery profiles (Wood et al.,  
69 2013) may be limited by match duration (80 min compared to 60 min for  
70 International matches), small sample size ( $n=6$ ) and single match design as opposed

71 to that of a tournament, leading to an underestimation of the responses to  
72 tournament match-play. Recent reports of match demands have differed (Fox et al.,  
73 2013) to previous reports in elite players (Otago, 1983), as such recent rule changes  
74 (January 2016), intended to reduce stoppages and increase the speed and intensity  
75 of match-play, may have compromised the application of previous literature  
76 regarding the demands and responses to netball match-play. Limited information  
77 exists regarding the external loads of professional netball, (Bailey et al., 2017;  
78 Young et al., 2016) and no studies have examined the physiological demands and  
79 responses to either a single or multiple instances of International-standard netball  
80 match-play. A deeper understanding of the movement patterns, coupled with  
81 physiological demands, can allow effective training to be prescribed to optimise  
82 adaptation and performance, however this information is currently limited (Bailey  
83 et al., 2017). Therefore, the purpose of this study was to examine the physiological,  
84 neuromuscular, endocrine and perceptual responses to an International netball  
85 tournament as well as the physiological demands of International-standard netball.

## 86 **Methods**

87 This observational study examined the response to a netball tournament performed  
88 over three consecutive days. Matches commenced at 19:00, 15:00 and 15:00 h on  
89 days one to three, respectively. On the morning of each match (~07:30 h), and three  
90 days (approximately 62 h) after the final match (~07:30 h), scores for perceived  
91 well-being (adapted brief assessment of mood+; BAM+), and samples of whole  
92 blood (Creatine Kinase concentration; CK) and saliva (cortisol; C and testosterone:  
93 T concentrations) were collected, and countermovement jump testing performed.  
94 Match intensity was quantified using both internal (heart rate telemetry) and  
95 external (accelerometry) load metrics. Following the match, players individually

96 recorded session (sRPE: Foster et al., 2001) and differential ratings of perceived  
97 exertion (dRPE: Weston, Siegler, Bahnert, McBrien, & Lovell, 2015) using a  
98 numerically blinded CR100® scale via an Android tablet. These values were  
99 recorded during the cool down period, ~15 min after match-play.

100

101 Eleven female players (age:  $25 \pm 4$  years; mass:  $71.8 \pm 7.8$  kg; height:  $1.8 \pm 0.1$  m)  
102 from an International netball team were recruited. Players were assigned according  
103 to positions to goal-based ( $n=2$ , GS and GK) and mid-court ( $n=9$ , GD, WD, WA,  
104 C and GA) groups based on court movement restrictions. This study included an  
105 International tournament played at the end of the 2016 domestic season. As such,  
106 all players had competed weekly in the British Super League (highest netball league  
107 in Britain) and were engaged in full-time training (strength, speed, endurance and  
108 netball-specific training sessions four to six times per week) as part of their club's  
109 performance preparation program. Five players used no form of hormonal  
110 contraceptive and players were requested to self-monitor menstrual cycles and days  
111 of contraceptive consumption. Subsequent analyses revealed no bias in hormonal  
112 markers as a function of contraceptive use. This study was approved by the Swansea  
113 University ethics committee, players were informed of the benefits and risks of the  
114 investigation before signing informed consent forms and completing health  
115 screening and were made aware that all material would be anonymised. All  
116 mandatory health and safety procedures were complied with in completing this  
117 research study.

118

119 Players completed BAM+ which is correlated to high-intensity match activity, and  
120 is sensitive to physiological responses following elite team sport match-play

121 (Shearer et al., 2017). Using an Android tablet (Iconia One 7 B1-750, Taipei,  
122 Taiwan: Acer inc), a series of questions was answered with a 100 mm visual  
123 analogue scale anchored with “not at all [0]” and “extremely [100]”. An overall  
124 recovery score was generated by subtracting the mean score of negative related  
125 items from the mean score of the positively related questions using Equation 1:  
126 (Shearer et al., 2017).

127

128 Equation 1: (Alertness + sleep quality + confidence + motivation) /4 - (Anger +  
129 confusion + tension + depression + fatigue + muscle soreness)/6.

130

131 For salivary hormone analysis, players were instructed to avoid eating food or  
132 drinking fluids other than water after waking to avoid contaminating saliva samples.  
133 Prior to breakfast, a two ml sample of saliva was collected via passive drool  
134 (Crewther et al., 2013) into sterile containers, with samples subsequently stored at  
135 -70°C until assay. After thawing and centrifugation (2000 revolutions·min<sup>-1</sup> for 10  
136 min), the samples were analysed in duplicate for T and C using commercial kits  
137 (Salimetrics, LLC, State College, PA, USA). The minimum detection limit for the  
138 testosterone assay was 6.1 pg·ml<sup>-1</sup>, with interassay coefficient of variation (CV)  
139 <10%. The cortisol assay had a detection limit of 0.12 ng·ml<sup>-1</sup> with interassay CV  
140 <7%. Samples for each player were assayed in the same plate to eliminate inter-  
141 assay variability.

142

143 Whole blood CK concentrations were measured via capillary blood (120 µl) being  
144 sampled from the fingertip and stored on ice in EDTA prepared collection tubes  
145 (Microvette 500, Sarstedt, Numbrecht, Germany) before being centrifuged at 3000

146 revolutions·min<sup>-1</sup> for 10 min (Labofuge 400R; Kendro Laboratories,  
147 Langensfeld, Germany). Plasma samples were then stored at -70°C before being  
148 analysed for CK concentration using commercially available kits (CK-NAC, ABX  
149 Diagnostics, Northampton, United Kingdom) on a spectrophotometer (Cobas Mira,  
150 ABX Diagnostics, Northampton, United Kingdom). Samples were measured in  
151 duplicate (CV=3%) and recorded as a mean.

152

153 A portable force platform with built-in charge amplifier (Kistler type 92866AA,  
154 Kistler Instruments Ltd., Farnborough, UK) measured the ground reaction force-  
155 time history of countermovement jumps. A sample rate of 1000 Hz was used, and  
156 the platform's calibration was confirmed prior to testing. Power (CV=2.4%) and  
157 jump height (JH; calculated from takeoff velocity; CV=3.4%) was calculated using  
158 previously established procedures (Owen et al., 2014; West et al., 2011) and have  
159 been reported to be sensitive to changes following competitive matches (Russell et  
160 al., 2015; West et al., 2014). Players performed a standardised warm up before  
161 jumping, placed hands on hips throughout the jump, and performed two jumps at  
162 each time-point with the best jump taken as the highest peak power output (PPO)  
163 and used in subsequent analyses.

164

165 External load was quantified by use of a microtechnology unit (Catapult S5,  
166 Catapult, Innovations, Leeds, UK) housing an in-built tri-axial accelerometer  
167 sampling at 100 Hz. Players wore a custom-made vest (Catapult Innovations,  
168 Leeds, UK) in which units were held in place vertically on the upper back to  
169 minimise movement. Data were downloaded using the manufacturer's software  
170 (Catapult sprint 5.1, Catapult Innovations, Leeds, UK), analysed for player-load for



171 each quarter, excluding breaks between quarters, with data represented as external  
172 load intensity ( $\text{AU}\cdot\text{min}^{-1}$ ). Data was pooled and reported for each position rather  
173 than individual players, such that for every match each position would have a single  
174 external load intensity for each quarter. Player-load has been reported to be a valid  
175 and reliable method (Barrett, Midgley, & Lovell, 2014; Boyd, Ball, & Aughey,  
176 2011) of measuring activities performed in team sports movements, with high  
177 within and between-device ( $\text{CV}\sim 1\%$ : Boyd et al., 2011) reliability and has been  
178 widely used in team sports (Luteberget & Spencer, 2017; Polgaze, Dawson,  
179 Hiscock, & Peeling, 2015) including netball (Chandler, Pinder, Curran, & Gabbett,  
180 2014; Young et al., 2016) with detailed calculations described previously (Barrett  
181 et al., 2014). Players wore heart rate (HR) monitors (Polar Team System 2, Polar  
182 Electro, Warwick, UK) throughout matches, with HR recorded at beat-to-beat  
183 intervals. Data was downloaded and analysed for each quarter, excluding breaks  
184 between quarters, and only whilst the player was on-court, using the Polar team  
185 system software (Polar Team 2, Polar Electro, Warwick, UK). HR data was  
186 reported for each player and associated to the position which had been played.

187

188 Following each match, players recorded sRPE along with indices of dRPE  
189 including ratings for breathlessness (RPE-B), leg muscle exertion (RPE-L), upper  
190 body muscle exertion (RPE-U) and cognitive/technical demands (RPE-T) (Weston  
191 et al., 2015). Ratings were provided using a numerically blinded CR100® scale  
192 with verbal anchors using a bespoke application on an Android tablet. dRPE  
193 provides a detailed quantification of internal load during team sport activities  
194 (McLaren, Smith, Spears, & Weston, 2017), is a sensitive marker of match exertion  
195 (Weston et al., 2015) and distinguishes between different areas of effort (McLaren

196 et al., 2017; Weston et al., 2015). Players must have performed a minimum of one  
197 quarter for sRPE and dRPE to be included in subsequent analyses.

198

199 Data are reported as mean difference  $\pm$  90% confidence limits unless otherwise  
200 stated. Visual inspection of the residual plots revealed evidence of  
201 heteroscedasticity; therefore, except for sRPE, dRPE, BAM+ and HR, analyses  
202 were performed on log transformed data. Separate mixed linear mixed models  
203 (SPSS v.24, Armonk, NY: IBM Corp) were used to examine the effect of  
204 tournament match-play on measures of physical exertion (external load, HR, sRPE,  
205 dRPE) and, thereafter, the effect of playing position on match physical exertion,  
206 and, the effects of tournament match-play on the players' neuromuscular,  
207 physiological and perceptual responses (PPO, JH, CK, T, C). In these models,  
208 match (match 1, match 2, match 3), playing position (mid-court, goal-based) and  
209 time (day 1, day 2, day 3, 3 days post), respectively were entered as the fixed effects.  
210 In all models, players were included as a random effect with random intercept to  
211 account for the dependency that arises from a hierarchical data structure such as  
212 ours (i.e., repeated measurements from the same players). From here, a custom-  
213 made spreadsheet (Hopkins, 2007) was used to determine magnitude based  
214 inferences (Batterham & Hopkins, 2006) for all differences, with inferences based  
215 on standardised thresholds for small, moderate, large and very large differences of  
216 0.2, 0.6, 1.2 and 2.0 of the pooled between-subject standard deviations (SD)  
217 (Hopkins, Marshall, Batterham, & Hanin, 2009). The chance of the difference being  
218 substantial or trivial was interpreted using the following scale: 25–75%, possibly;  
219 75–95%, likely; 95–99.5%, very likely; >99.5%, most likely (Batterham &  
220 Hopkins, 2006). Uncertainty in all estimates is expressed via 90% confidence limits

221 and the magnitude of effects assessed mechanistically, whereby if the confidence  
222 limits overlapped the thresholds for the smallest worthwhile positive and negative  
223 effects, effects were deemed unclear (Hopkins et al., 2009).

## 224 **Results**

225 Match data are presented in Table 1. Mean playing time for players across the three  
226 matches was 119.8 min ( $\pm 48.5$  min;  $\pm$  SD) and outcomes included two wins and a  
227 loss for matches one to three respectively. In response to a single netball match,  
228 from day one to day two, CK (likely very large;  $72.6\% \pm 26.4\%$ ) and fatigue (likely  
229 small;  $56.2\% \pm 46.0\%$ ) increased, whilst motivation (likely moderate;  $-19.5\% \pm$   
230  $14.3\%$ ), BAM+ (likely moderate;  $-27.9\% \pm 17.6\%$ ), sleep quality (possibly  
231 moderate;  $-16.3\% \pm 15.6\%$ ), C (likely small;  $-27.4\% \pm 23.7\%$ ), T (possibly small;  
232  $-10.8\% \pm 10.8\%$ ) and JH (possibly small;  $-4.0\% \pm 2.5\%$ ) decreased, with a possible  
233 trivial difference for PPO and unclear difference for soreness (Table 2). Following  
234 two netball matches, from day one to day three, CK (most likely very large;  $120.8\%$   
235  $\pm 33.7\%$ ), fatigue (possibly large;  $146.9\% \pm 46.0\%$ ) and soreness (possibly  
236 moderate;  $57.7\% \pm 37.9\%$ ) increased, whilst BAM+ (likely large;  $-42.8 \pm 17.6\%$ ),  
237 motivation (likely moderate;  $-20.6\% \pm 14.3\%$ ), sleep quality (possibly moderate;  $-$   
238  $30.8\% \pm 15.6\%$ ), T (possibly small;  $-8.7\% \pm 11.0\%$ ) and PPO (possibly small;  $-$   
239  $3.3\% \pm 1.7\%$ ) decreased, with a possible trivial difference for JH and most likely  
240 trivial difference for C. Following the performance of two consecutive matches,  
241 from day two to three, CK (likely moderate;  $27.9\% \pm 19.5\%$ ), fatigue (likely  
242 moderate;  $58.1\% \pm 29.5\%$ ), soreness (possible moderate;  $49.6\% \pm 36.0\%$ ) and C  
243 (possibly moderate;  $43.3\% \pm 46.8\%$ ) increased whilst BAM+ (possibly moderate;  
244  $-20.6\% \pm 24.4\%$ ) and sleep quality (possibly moderate;  $-17.3\% \pm 18.6\%$ ) decreased,  
245 with an unclear difference for T and motivation, and likely trivial difference for JH

246 and PPO. Three days post-tournament BAM+ (likely very large;  $-57.5\% \pm 20.5\%$ ),  
247 sleep quality (likely large;  $-38.7\% \pm 18.1\%$ ), motivation (likely moderate;  $-24.3\%$   
248  $\pm 16.6\%$ ), PPO (likely small;  $-4.2\% \pm 1.9\%$ ), JH (possibly small;  $-3.9\% \pm 2.8\%$ )  
249 and T (possibly small;  $-10.0\% \pm 12.7\%$ ) decreased, whilst fatigue increased (very  
250 likely moderate;  $127.2\% \pm 53.6\%$ ) compared to day one, with unclear differences  
251 for C, CK and soreness.

252

253 *\*\*\*\*Table 1 about here\*\*\*\**

254 *\*\*\*\*Table 2 about here\*\*\*\**

255

256 Greater mean HR for match one occurred relative to match two (possibly small;  
257  $1.2\% \pm 0.02\%$ ). Likely trivial differences were observed for external load intensity  
258 and unclear differences for sRPE and dRPE variables. For match three versus one  
259 for RPE-B (likely small;  $20.1\% \pm 25.4\%$ ), RPE-L (possibly small;  $18.2\% \pm 24.5\%$ ),  
260 RPE-U (possibly small;  $18.1\% \pm 22.4\%$ ) and RPE-T (possibly moderate;  $23.2\% \pm$   
261  $19.8\%$ ), greater values were observed. A possible trivial difference existed for  
262 external load intensity and unclear differences for sRPE and mean HR. Match three  
263 produced greater sRPE (likely small;  $21.7\% \pm 27.4\%$ ), RPE-B (possibly moderate;  
264  $32.0\% \pm 26.7\%$ ), RPE-L (possibly moderate;  $30.8\% \pm 25.9\%$ ), RPE-U (likely small;  
265  $30.6\% \pm 23.7\%$ ), RPE-T (possibly moderate;  $27.1\% \pm 20.2\%$ ) and mean HR  
266 (possibly small;  $1.1\% \pm 2.0\%$ ) versus match two. There was a possible trivial  
267 difference for external load intensity.

268

269 Overall, mid-court positions performed at a greater external load intensity (possibly  
270 very large;  $85.7\% \pm 49.6\%$ ), mean HR (possibly moderate;  $3.7\% \pm 3.8\%$ ) (Table

271 3), and reported higher sRPE (possibly moderate;  $40.7\% \pm 40.0\%$ ), RPE-B (likely  
272 moderate;  $55.9\% \pm 51.9\%$ ), RPE-L (possibly large;  $79.3\% \pm 48.1\%$ ), RPE-U  
273 (possibly moderate;  $47.2\% \pm 54.9\%$ ) and RPE-T (possibly moderate;  $36.9\% \pm$   
274  $36.7\%$ ) compared to goal-based positions (Table 4).

275

276 *\*\*\*\*Table 3 about here\*\*\*\**

277 *\*\*\*\*Table 4 about here\*\*\*\**

## 278 **Discussion**

279 The aims of this study were to characterise the physiological, neuromuscular,  
280 endocrine and perceptual responses to an International tournament and to identify  
281 the position-specific demands of International netball. The primary findings were  
282 that the performance of both a single, and multiple matches resulted in a varied  
283 recovery profile, with greater perturbations in perceived well-being and  
284 physiological function following consecutive matches, and fatigue evident up to  
285 three days post-tournament. Additionally, mid-court positions performed at greater  
286 internal and external load intensity compared to goal-based positions.

287

288 Across the tournament, CK, reported to be indicative of skeletal muscle damage  
289 (Cunniffe et al., 2010), accumulated before returning to baseline thereafter. Whilst  
290 there are no reports in netball, investigations in other team sports have reported  
291 peak values occurring 24 h post-match, remaining elevated for females for up to 69  
292 h (Andersson et al., 2008). Three days post-tournament, CK and perceived soreness  
293 had returned to baseline, however neuromuscular performance and T  
294 concentrations remained suppressed. This may suggest that neuromuscular  
295 performance is impacted by T concentration rather than muscle damage, that CK is

296 not sensitive to detect changes in muscle damage, or that various markers of fatigue  
297 collectively interact.

298

299 Following the performance of a single match, T was reduced, and remained  
300 reduced until three days post-tournament, whilst C decreased following the first  
301 match, then returned and remained at baseline following the second match.

302 Testosterone concentration is associated with enhanced neuromuscular  
303 performance (Cook, Kilduff, Crewther, Beaven, & West, 2014), decision making,  
304 behaviour, contractile signalling (Crewther, Cook, Cardinale, & Weatherby, 2011),  
305 motivation (Cook, Kilduff, & Crewther, 2018) and performance (Crewther et al.,  
306 2013). A reduction, as seen in the present study, may have negatively affected one  
307 or more of these reported associations, with a resultant impact upon performance.

308 The recovery of C following two matches may suggest a varied anticipatory  
309 response with a greater anticipatory rise prior to the first and final match (higher  
310 ranked opponent for the final match). However, alternatively the late  
311 commencement (19:00 h compared to 15:00 h) of match one may have negatively  
312 affected post-match processes and recovery. Menstrual phase and hormonal  
313 contraceptive use were not controlled for in the present study, however no  
314 difference was found in basal T between hormonal contraceptive users and non-  
315 users. Additionally, recent reports highlight only a difference in magnitude of T  
316 response to a stimulus, rather than the response itself, and no impact upon  
317 performance with hormonal contraceptive use (Cook et al., 2018).

318

319 This is the first study to characterise playing demands during an International  
320 tournament reporting external load, perceived effort and HR. Internal and external

321 load was greater for mid-court compared to goal-based positions (Table 3). Greater  
322 external load intensity for mid-court positions has been previously reported in  
323 professional netball (Fox et al., 2013; Young et al., 2016), and is likely due to court  
324 movement restrictions resulting in a higher active time (Fox et al., 2013), time spent  
325 in high-intensity zones (Young et al., 2016) and type of on and off-ball locomotor  
326 and non-locomotor activity (Bailey et al., 2017). Collectively, this suggests that  
327 players should not only be conditioned for the position specific movement  
328 demands, as previously reported, but also the different physiological and type of  
329 effort (as indicated by dRPE) experienced during International match-play. Both  
330 sRPE and dRPE can be used by conditioning staff to guide the individualisation of  
331 the training stimulus to the positional demands. As markers of fatigue were further  
332 reduced following a greater number of consecutive matches, training should aim to  
333 replicate these demands to minimise this disturbance, especially when considering  
334 that some International tournaments are up to twice as long as in the present study.  
335 Unlike perceptual and endocrine responses, neuromuscular performance was not  
336 further reduced following consecutive matches. Perceptual markers could therefore  
337 be considered as a simple monitoring tool to identify sufficient training load to  
338 replicate the fatiguing consequences associated with International netball. Sleep  
339 quality was negatively affected following a single, and to a greater extent following  
340 consecutive matches, a consideration for coaching and support staff, as sleep has  
341 been reported to be vital for recovery (Halson, 2008). Three days post-tournament,  
342 when players commenced training, perceived well-being, sleep quality, T  
343 concentration and neuromuscular function were reduced, suggesting longer  
344 recovery is required than anticipated by conditioning staff.

345 **Conclusion**

346 This is the first study to report the physiological demands of and responses to an  
347 International netball tournament, providing vital information for International  
348 coaches and conditioning coaches. Markers of fatigue increased following the  
349 performance of a single match, whilst markers of muscle damage and perceived  
350 well-being were further affected following consecutive matches. A varied recovery  
351 profile was apparent as recovery to baseline of all variables examined did not occur  
352 62 h post-tournament. Mid-court positions performed at higher external and internal  
353 intensities compared to goal-based positions, an important consideration for  
354 conditioning staff in order to individualise training to positional specific demands.

355

356

357

358

359

360

361 **Acknowledgements**

362 We would like to acknowledge the Welsh netball players, coaching and support staff for  
363 their support through this testing. Additionally, to Liam Harper, Christopher Lewis  
364 and Charlie Finn for assisting with data collection.

365

366 This research did not receive any specific grant from funding agencies in the public,  
367 commercial, or not-for-profit sectors.

368

369 **Disclosure of interest**

370 The authors report no conflict of interest.

371

372

373



374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396

397 **References**

- 398 Andersson, H., Raastad, T., Nilsson, J., Paulsen, G., Garthe, I., & Kadi, F. (2008).  
399 Neuromuscular fatigue and recovery in elite female soccer: Effects of active  
400 recovery. *Medicine and Science in Sports and Exercise*, 40(2), 372–380.  
401 <https://doi.org/10.1249/mss.0b013e31815b8497>
- 402 Bailey, J. A., Gastin, P. B., Mackey, L., & Dwyer, D. B. (2017). The Player Load  
403 Associated With Typical Activities in Elite Netball. *International Journal of*  
404 *Sports Physiology and Performance*, 12, 1218–1223.  
405 <https://doi.org/10.1123/ijsspp.2016-0378>

- 406 Barrett, S., Midgley, A., & Lovell, R. (2014). PlayerLoad<sup>TM</sup>: Reliability,  
407 convergent validity, and influence of unit position during treadmill running.  
408 *International Journal of Sports Physiology and Performance*, 9(6), 945–952.  
409 <https://doi.org/10.1123/ijssp.2013-0418>
- 410 Batterham, A. M., & Hopkins, W. G. (2006). Making meaningful inferences about  
411 magnitudes. *International Journal of Sport Physiology and Performance*,  
412 1(1), 50–57.
- 413 Boyd, L. J., Ball, K., & Aughey, R. J. (2011). The reliability of MinimaxX  
414 accelerometers for measuring physical activity in Australian football.  
415 *International Journal of Sport Physiology and Performance*, 6, 311–321.  
416 <https://doi.org/10.1123/ijssp.6.3.311>
- 417 Chandler, P., Pinder, S., Curran, J., & Gabbett, T. J. (2014). Physical demands of  
418 training and competition in collegiate netball players, 28(10), 2732–2737.  
419 <https://doi.org/10.1519/JSC.0000000000000486>
- 420 Cook, C. J., Kilduff, L. P., & Crewther, B. T. (2018). Basal and stress-induced  
421 salivary testosterone variation across the menstrual cycle and linkage to  
422 motivation and muscle power. *Scandinavian Journal of Medicine & Science  
423 in Sports*, 28(4), 1345–1353. <https://doi.org/10.1111/sms.13041>
- 424 Cook, C. J., Kilduff, L. P., Crewther, B. T., Beaven, M., & West, D. J. (2014).  
425 Morning based strength training improves afternoon physical performance in  
426 rugby union players. *Journal of Science and Medicine in Sport*, 17(3), 317–  
427 321. <https://doi.org/10.1016/j.jsams.2013.04.016>
- 428 Crewther, B. T., Cook, C., Cardinale, M., & Weatherby, R. (2011). Two emerging  
429 concepts for elite athletes cortisol and testosterone. *Review Literature And  
430 Arts Of The Americas*, 41(2), 103–123. <https://doi.org/10.2165/11536850->

431 000000000-00000

432 Crewther, B. T., Sanctuary, C. E., Kilduff, L. P., Carruthers, J. S., Gaviglio, C.  
433 M., & Cook, C. J. (2013). The Workout Responses of Salivary-Free  
434 Testosterone and Cortisol Concentrations and Their Association With the  
435 Subsequent Competition Outcomes in Professional Rugby League. *Journal*  
436 *of Strength and Conditioning Research*, 27(2), 471–476.  
437 <https://doi.org/10.1519/JSC.0b013e3182577010>

438 Cunniffe, B., Hore, A. J., Whitcombe, D. M., Jones, K. P., Baker, J. S., & Davies,  
439 B. (2010). Time course of changes in immunoendocrine markers following  
440 an international rugby game. *European Journal of Applied Physiology*,  
441 108(1), 113–122. <https://doi.org/10.1007/s00421-009-1200-9>

442 Foster, C., Florhaug, J., Franklin, J., Gottschall, L., Hrovatin, L., Parker, S., ...  
443 Dodge, C. (2001). A new approach to monitoring exercise training. *Journal*  
444 *of Strength and Conditioning Research*, 15(1), 109–115.

445 Fox, A., Spittle, M., Otago, L., & Saunders, N. (2013). Activity profiles of the  
446 Australian female netball team players during international competition:  
447 implications for training practice. *Journal of Sports Sciences*, 31(14), 1588–  
448 1595. <https://doi.org/10.1080/02640414.2013.792943>

449 Halson, S. L. (2008). Nutrition, sleep and recovery. *European Journal of Sport*  
450 *Science*, 8(2), 119–126. <https://doi.org/10.1080/17461390801954794>

451 Hopkins, W. G. (2007). A spreadsheet for deriving confidence interval,  
452 mechanistic inference and clinical inference from a P value. *SportScience*,  
453 11, 16+.

454 Hopkins, W. G., Marshall, S. W., Batterham, A. M., & Hanin, J. (2009).  
455 Progressive statistics for studies in sports medicine and exercise science.

456 *Medicine and Science in Sports and Exercise*, 41(1), 3–12.  
457 <https://doi.org/10.1249/MSS.0b013e31818cb278>

458 Luteberget, L. S., & Spencer, M. (2017). High-intensity events in international  
459 women's team handball matches. *International Journal of Sports Physiology*  
460 *and Performance*, 12(1), 56–61. <https://doi.org/10.1123/ijsp.2015-0641>

461 McLaren, S. J., Smith, A., Spears, I. R., & Weston, M. (2017). A detailed  
462 quantification of differential ratings of perceived exertion during team-sport  
463 training. *Journal of Science and Medicine in Sport*, 20(3), 290–295.  
464 <https://doi.org/10.1016/j.jsams.2016.06.011>

465 Otago, L. (1983). A game analysis of the activity patterns of netball players.  
466 *Sports Coach*, 7, 24–28.

467 Owen, N., Watkins, J., Kilduff, L., Cunningham, D., Bennett, M., & Bevan, H.  
468 (2014). Development of a criterion methods to determine peak mechanical  
469 power output in a countermovement jump. *Journal of Strength and*  
470 *Conditioning Research*, 28(6), 1552–1558.

471 Polgaze, T., Dawson, B., Hiscock, D., & Peeling, P. (2015). A comparative  
472 analysis of accelerometer and time motion data in elite mens hockey training  
473 and competition. *International Journal of Sports Physiology and*  
474 *Performance*, 10(4), 446–451. <https://doi.org/10.1123/ijsp.2014-0233>

475 Russell, M., Northeast, J., Atkinson, G., Shearer, D. A., Sparkes, W., Cook, C. J.,  
476 & Kilduff, L. P. (2015). Between-Match Variability of Peak Power Output  
477 and Creatine Kinase Responses to Soccer Match-Play. *Journal of Strength*  
478 *and Conditioning Research*, 29(8), 2079–2085.  
479 <https://doi.org/10.1519/JSC.0000000000000852>

480 Shearer, D. A., Sparkes, W., Northeast, J., Cunningham, D. J., Cook, C. J., &

481 Kilduff, L. P. (2017). Measuring recovery: An adapted Brief Assessment of  
482 Mood (BAM+) compared to biochemical and power output alterations.  
483 *Journal of Science and Medicine in Sport*, 20(5), 512–517.  
484 <https://doi.org/10.1016/j.jsams.2016.09.012>

485 West, D. J., Cook, C. J., Stokes, K. A., Atkinson, P., Drawer, S., Bracken, R. M.,  
486 & Kilduff, L. P. (2014). Profiling the time-course changes in neuromuscular  
487 function and muscle damage over two consecutive tournament stages in elite  
488 rugby sevens players. *Journal of Science and Medicine in Sport*, 17(6), 688–  
489 692. <https://doi.org/10.1016/j.jsams.2013.11.003>

490 West, D. J., Owen, N. J., Jones, M. R., Bracken, R. M., Cook, C. J., Cunningham,  
491 D. J., ... Kilduff, L. P. (2011). Relationships between force-time  
492 characteristics of the isometric midthigh pull and dynamic performance in  
493 professional rugby league players. *Journal of Strength and Conditioning*  
494 *Research*, 25(11), 3070–3075.  
495 <https://doi.org/10.1519/JSC.0b013e318212dcd5>

496 Weston, M., Siegler, J., Bahnert, A., McBrien, J., & Lovell, R. (2015). The  
497 application of differential ratings of perceived exertion to Australian Football  
498 League matches. *Journal of Science and Medicine in Sport*, 18(6), 704–708.  
499 <https://doi.org/10.1016/j.jsams.2014.09.001>

500 Wood, A., Kelly, V., & Gabbett, T. J. (2013). Neuromuscular And Perceptual  
501 Fatigue Responses To An Elite Level Netball Match. *Journal of Australian*  
502 *Strength and Conditioning*, 21(1), 24–29.

503 Young, C. M., Gatin, P. B., Sanders, N., Mackey, L., & Dwyer, D. B. (2016).  
504 Player load in elite netball: Match, training, and positional comparisons.  
505 *International Journal of Sports Physiology and Performance*, 11(8), 1074–

506

1079. <https://doi.org/10.1123/ijsp.2015-0156>

507

508 **Table 1:** Mean  $\pm$  SD heart rate (absolute and percent of age predicted maximum),  
 509 sRPE, indices of dRPE and external load intensity for each match averaged across  
 510 all players ( $n=31$ ). Magnitude of difference and uncertainty shown between each  
 511 match.

	Match 1	1 v 2	Match 2	2 v 3	Match 3	3 v 1	Overall
Mean HR (b·min <sup>-1</sup> )	170 ( $\pm 8.0$ )	S*	169 ( $\pm 9.4$ )	S*	171 ( $\pm 9.2$ )	U	170 ( $\pm 8.7$ )
Mean HR (% of max)	87.7 ( $\pm 3.6$ )	S*	86.7 ( $\pm 4.4$ )	S*	88.0 ( $\pm 4.8$ )	U	87.5 ( $\pm 4.2$ )
sRPE (AU)	63.4 ( $\pm 20.3$ )	U	58.6 ( $\pm 30.2$ )	S**	73.0 ( $\pm 29.3$ )	U	65.3 ( $\pm 26.9$ )
RPE-B (AU)	56.2 ( $\pm 27.1$ )	U	51.5 ( $\pm 28.1$ )	M*	71.5 ( $\pm 34.7$ )	S**	60.1 ( $\pm 30.6$ )
RPE-L (AU)	57.1 ( $\pm 21.2$ )	U	52.1 ( $\pm 28.4$ )	M*	70.7 ( $\pm 30.6$ )	S*	60.3 ( $\pm 27.5$ )
RPE-U (AU)	38.1 ( $\pm 22.2$ )	U	35.9 ( $\pm 25.7$ )	S**	50.3 ( $\pm 25.3$ )	S*	41.7 ( $\pm 24.5$ )
RPE-T (AU)	53.3 ( $\pm 19.5$ )	U	51.5 ( $\pm 19.1$ )	M*	67.7 ( $\pm 25.9$ )	M*	57.8 ( $\pm 22.4$ )
External load (AU·min <sup>-1</sup> )	7.9 ( $\pm 1.9$ )	T**	8.2 ( $\pm 2.3$ )	T*	8.7 ( $\pm 2.6$ )	T*	8.2 ( $\pm 2.2$ )

512 *Magnitude of the difference:* U: unclear; T: trivial; S: small; M: moderate; L: large;  
 513 VL: very large. *Uncertainty of the difference:* \*: possibly (25-75% (likelihood of  
 514 the difference being...); \*\*: likely (75-95%); \*\*\*: very likely (95-99.5%); \*\*\*\*:  
 515 most likely (>99.5%). *Abbreviations:* AU: arbitrary unit; SD: standard deviation;  
 516 sRPE: session rating of perceived exertion; dRPE: differential rating of perceived  
 517 exertion; RPE-B: rating of perceived breathlessness; RPE-L: rating of perceived leg  
 518 muscle exertion; RPE-U: rating of perceived upper body muscle exertion; RPE-T:  
 519 rating of perceived cognitive/ technical demand.

520

521 **Table 2:** Mean  $\pm$  SD cortisol, testosterone and CK concentrations, jump height,  
 522 peak power output and perceived well-being across the three days of the tournament  
 523 and three days post. Magnitude and uncertainty of the difference shown compared  
 524 to day one.

	Day 1	Day 2	Day 3	3 days post
Cortisol ( $\mu\text{g}\cdot\text{dl}^{-1}$ )	0.61 ( $\pm 0.25$ )	0.47 ( $\pm 0.23$ ) S**	0.65 ( $\pm 0.29$ ) T*****	0.58 ( $\pm 0.34$ ) U
Testosterone ( $\text{pg}\cdot\text{ml}^{-1}$ )	116.2 ( $\pm 33.5$ )	102.9 ( $\pm 25.9$ ) S*	105.4 ( $\pm 25.3$ ) S*	95.7 ( $\pm 27.0$ ) S*
CK ( $\text{U}\cdot\text{L}^{-1}$ )	123.3 ( $\pm 30.9$ )	217.2 ( $\pm 67.4$ ) VL**	283.0 ( $\pm 121.3$ ) VL*****	141.9 ( $\pm 113.0$ ) U
PPO (W)	3311 ( $\pm 440$ )	3235 ( $\pm 389$ ) T*	3194 ( $\pm 369$ ) S*	3120 ( $\pm 294$ ) S**
Jump height (m)	0.30 ( $\pm 0.05$ )	0.29 ( $\pm 0.04$ ) S*	0.29 ( $\pm 0.04$ ) T*	0.30 ( $\pm 0.16$ ) S*
BAM+ (AU)	51.5 ( $\pm 15.2$ )	37.2 ( $\pm 21.7$ ) M**	29.6 ( $\pm 20.2$ ) L**	23.6 ( $\pm 30.6$ ) VL**
Soreness (AU)	31.8 ( $\pm 23.6$ )	33.5 ( $\pm 21.4$ ) U	50.2 ( $\pm 20.5$ ) M*	41.6 ( $\pm 25.6$ ) U
Fatigue (AU)	17.6 ( $\pm 19.0$ )	27.5 ( $\pm 9.1$ ) L*	43.5 ( $\pm 15.8$ ) L*	42.3 ( $\pm 20.1$ ) M***
Sleep quality (AU)	76.5 ( $\pm 18.0$ )	64.1 ( $\pm 24.8$ ) M*	53.0 ( $\pm 24.6$ ) M*	48.1 ( $\pm 24.9$ ) L**
Motivation (AU)	75.5 ( $\pm 15.5$ )	60.7 ( $\pm 25.7$ ) M**	59.9 ( $\pm 18.2$ ) M**	60.0 ( $\pm 16.5$ ) M**

525 *Magnitude of the difference:* U: unclear T: trivial; S: small; M: moderate; L: large;  
 526 VL: very large. *Uncertainty of the difference:* \*: possibly (25-75% (likelihood of  
 527 the difference being...); \*\*: likely (75-95%); \*\*\*: very likely (95-99.5%); \*\*\*\*\*:  
 528 most likely (>99.5%). *Abbreviations:* AU: arbitrary unit; SD: standard deviation;  
 529 CK: creatine kinase concentration; PPO: peak power output; BAM+: adapted brief  
 530 assessment of mood.



531 **Table 3:** Mean  $\pm$  SD heart rate (absolute and percent of age predicted maximum) and external load intensity for each match and averaged across  
 532 all matches for mid-court and goal-based positional groups. Magnitude of difference and uncertainty shown between positional groups.

		Match 1	Match 2	Match 3	Mean	Difference between positional groups
<i>Mid-court (n=24)</i>	Mean HR (b·min <sup>-1</sup> )	170 ( $\pm$ 8.9)	172 ( $\pm$ 8.7)	173 ( $\pm$ 7.2)	172 ( $\pm$ 7.9)	M*
	Mean HR (% of max)	88.1 ( $\pm$ 4.0)	88.6 ( $\pm$ 3.4)	89.1 ( $\pm$ 3.1)	88.6 ( $\pm$ 3.4)	M*
<i>Mid-court (n=15)</i>	External load (AU·min <sup>-1</sup> )	8.9 ( $\pm$ 0.8)	9.4 ( $\pm$ 0.8)	10.0 ( $\pm$ 0.6)	9.4 ( $\pm$ 0.8)	VL*
<i>Goal-based (n=7)</i>	Mean HR (b·min <sup>-1</sup> )	168 ( $\pm$ 4.0)	160 ( $\pm$ 4.5)	162 ( $\pm$ 14.6)	162 ( $\pm$ 7.6)	
	Mean HR (% of max)	86.1 ( $\pm$ 0.5)	82.5 ( $\pm$ 3.8)	83.1 ( $\pm$ 9.9)	83.7 ( $\pm$ 4.9)	
<i>Goal-based (n=6)</i>	External load (AU·min <sup>-1</sup> )	5.2 ( $\pm$ 0.5)	5.2 ( $\pm$ 1.8)	5.3 ( $\pm$ 2.6)	5.2 ( $\pm$ 1.4)	

533 *Magnitude of the difference:* U: unclear T: trivial; S: small; M: moderate; L: large; VL: very large. *Uncertainty of the difference:* \*: possibly (25-  
 534 75% (likelihood of the difference being...); \*\*: likely (75-95%); \*\*\*: very likely (95-99.5%); \*\*\*\*: most likely (>99.5%). *Abbreviations:* SD:  
 535 standard deviation; HR: heart rate; AU: arbitrary unit.

536

537

538

539 **Table 4:** Mean  $\pm$  SD sRPE and dRPE for each match and averaged across all matches for mid-court and goal-based positional groups. Magnitude  
 540 of difference and uncertainty shown between positional groups.

		Match 1	Match 2	Match 3	Mean	Difference between positional groups
<i>Mid-court (n=24)</i>	sRPE (AU)	64.9 ( $\pm$ 20.7)	66.3 ( $\pm$ 33.6)	76.0 ( $\pm$ 28.6)	69.5 ( $\pm$ 27.1)	M*
	RPE-B (AU)	60.4 ( $\pm$ 28.8)	57.3 ( $\pm$ 32.3)	74.4 ( $\pm$ 35.2)	64.8 ( $\pm$ 31.9)	M**
	RPE-L (AU)	60.8 ( $\pm$ 22.2)	59.4 ( $\pm$ 30.7)	76.3 ( $\pm$ 28.9)	66.2 ( $\pm$ 27.4)	L*
	RPE-U (AU)	42.0 ( $\pm$ 23.3)	42.1 ( $\pm$ 28.9)	53.9 ( $\pm$ 25.7)	46.5 ( $\pm$ 25.5)	M*
	RPE-T (AU)	59.8 ( $\pm$ 15.8)	56.4 ( $\pm$ 17.2)	68.2 ( $\pm$ 27.2)	62.0 ( $\pm$ 20.9)	M*
<i>Goal-based (n=7)</i>	sRPE (AU)	57.5 ( $\pm$ 24.7)	40.7 ( $\pm$ 6.5)	59.5 ( $\pm$ 40.3)	50.9 ( $\pm$ 21.9)	
	RPE-B (AU)	39.5 ( $\pm$ 10.6)	38.0 ( $\pm$ 5.3)	58.0 ( $\pm$ 41.0)	44.1 ( $\pm$ 20.0)	
	RPE-L (AU)	42.5 ( $\pm$ 9.2)	35.0 ( $\pm$ 13.7)	45.5 ( $\pm$ 33.2)	40.1 ( $\pm$ 16.9)	
	RPE-U (AU)	22.5 ( $\pm$ 3.5)	21.3 ( $\pm$ 4.7)	34.0 ( $\pm$ 21.2)	25.3 ( $\pm$ 11.0)	
	RPE-T (AU)	27.5 ( $\pm$ 0.7)	39.3 ( $\pm$ 21.0)	65.5 ( $\pm$ 27.6)	43.3 ( $\pm$ 23.0)	

541 *Magnitude of the difference:* U: unclear T: trivial; S: small; M: moderate; L: large; VL: very large. *Uncertainty of the difference:* \*: possibly (25-  
 542 75% (likelihood of the difference being...); \*\*: likely (75-95%); \*\*\*: very likely (95-99.5%); \*\*\*\*: most likely (>99.5%). *Abbreviations:* AU:  
 543 arbitrary unit; SD: standard deviation; sRPE: session rating of perceived exertion; dRPE: differential rating of perceived exertion; RPE-B: rating  
 544 of perceived breathlessness; RPE-L: rating of perceived leg muscle exertion; RPE-U: rating of perceived upper body muscle exertion; RPE-T:  
 545 rating of perceived cognitive/ technical demand.